




Introduction

BodyZone Physiotherapy for Ankle



Welcome to BodyZone Physiotherapy's patient resource about Peroneal Tendon Subluxation.

The peroneals are two muscles and their tendons that attach along the outer edge of the lower leg. The peroneal tendons are enclosed in a fibrous tunnel that runs behind the outside ankle bone (the lateral malleolus). Damage or injury to the structures that form and support this tunnel may lead to a condition in which the peroneal tendons snap out of place. This condition is called peroneal tendon subluxation.

This article will help you understand:


- how peroneal tendon subluxation happens
- how doctors diagnose the condition

- what can be done to treat this problem

Anatomy

The primary muscles supporting the **lateral (outer) part of the ankle** are the peroneals. These two muscles and their tendons lie along the outside of the lower leg bone (*fibula*) and cross behind the lateral malleolus (the outside ankle bone).

Lateral Part of Ankle




The tendons of the peroneal muscles pass together through a groove behind the lateral malleolus. The tendons are held within the groove by a sheath that forms a tunnel around the tendons. The surface of this sheath is reinforced by a ligament called a *retinaculum*. Contracting the peroneals makes the tendons glide in the groove like a pulley. This action points the foot downward (*plantarflexion*) and outward (*eversion*).




Causes

Tendons attach muscles to bone. Tightening a muscle puts tension on the tendon, which can move bone. Many of the joints in the body are held in place by supportive connective tissue, such as a ligament or retinaculum. If the supportive tissue has been **damaged or injured**, the tendon may be free to slip out of its normal position. This is called **subluxation**. When a subluxed tendon slips back into place, this is called *relocating*. A tendon that forcefully snaps out of position and does not relocate has *dislocated*.


Damaged Supportive Tissue



Subluxation



The main cause of peroneal tendon subluxation is an ankle sprain. A sprain that injures the ligaments on the outer side of the ankle can also damage the peroneal tendons. During the typical inversion ankle sprain, the foot rolls in. The force of the stretch on the peroneals can rip the retinaculum that keeps the peroneal tendons positioned in the groove. As a result, the tendons can jump out of the groove. The tendons usually relocate by snapping back into place.



The injury to the retinaculum may be overlooked at first while treatment focuses on the injury to other ankle ligaments. This means the subluxation may begin much later, and it may not seem to be caused by the initial ankle sprain. If not treated, this snapping of the tendons can become a chronic and recurring problem.

An acute dislocation of the peroneal tendons is rare. It occasionally happens during sport activities that force the foot to turn up and in, for example during skiing, ice skating, or soccer. At the moment the foot turns up and in, the peroneals violently contract to protect the ankle. This can cause the retinaculum to tear, allowing the tendons to slip out of the groove.

Differences in the anatomy of the groove may predispose some people to peroneal tendon subluxations. The groove may be too shallow. Or the ridge that helps deepen this groove may be too small or even absent. Sometimes, the retinaculum that keeps the tendons in the groove may be too loose. In these cases, patients may not recall any injury to explain the snapping of the peroneal tendons.

Symptoms

What does peroneal tendon subluxation feel like?

Patients describe a popping or snapping sensation on the outer edge of the ankle. The tendons may even be seen protruding out of place along the lower tip of the fibula. It is common to feel pain and tenderness **along the tendons**. There may be swelling just behind the bottom edge of the fibula.



Diagnosis

Diagnosis of peroneal subluxation begins with an examination of the ankle. Your physiotherapist at BodyZone Physiotherapy will move your ankle into different positions to see when the tendons snap out of place and if they return to place. One test involves holding pressure down on the ankle as you pull your foot up and out. Our physiotherapist feels for the fibula during this test to determine if the tendons are popping out of place. If your physiotherapist suspects a tear of the retinaculum, you may be referred for additional medical evaluation by a doctor.

Our Treatment

Non-surgical Rehabilitation

Nonsurgical treatment for peroneal tendon subluxations includes a program of rehabilitation exercises to help control symptoms. At BodyZone Physiotherapy, our physiotherapists will create a specialized program to help you improve strength and coordination of your ankle, and more quickly regain normal ankle function.

Post-surgical Rehabilitation

When you visit BodyZone Physiotherapy, your initial treatments will be used to help control pain and swelling following surgery. Our physiotherapist may apply ice and electrical stimulation during your first few physiotherapy sessions. We use massage and other hands-on treatments to ease muscle spasm and pain. Our physiotherapy is designed to help improve your range of motion without putting too much strain on the injured area.

Typically, after about six weeks of treatment, our therapist will help you increase your activity level. Your physiotherapist will slowly add exercises to improve the strength in your peroneal muscles and help you regain position sense in the ankle joint to improve its overall stability.

Rehabilitation after surgery can be a slow process, but our professionals will custom design a physiotherapy program for you.

speed your recovery. Recovery and rehabilitation varies for each patient, but as a guideline, you may expect to attend therapy sessions for two to three months, with full recovery typically requiring up to six months.

At BodyZone Physiotherapy, our goal is to help manage your pain, improve your range of motion, and maximize strength and control in your ankle. When your recovery is well under way, regular visits to our office will end. We will continue to be a supportive resource, but you will be in charge of doing your exercises as part of an ongoing home program.

Physician Review

If your doctor suspects a tear in the retinaculum, X-rays may be taken to determine if the torn retinaculum has pulled a piece of the fibula bone. This is called an avulsion fracture. X-rays are also used to look for other injuries to the ankle.

Your doctor may also order a magnetic resonance imaging (MRI) scan of your ankle. MRI scans can show abnormal swelling and scar tissue or tears in the tendons. However, MRIs won't always show subluxation of the peroneal tendons.

If your injury is acute, treatment without surgery may involve placing your ankle in a short-leg cast for four to six weeks. Your physician may also prescribe medications. Anti-inflammatories can help ease pain and swelling and get you back to normal activity sooner. These medications include common over-the-counter drugs such as ibuprofen. The goals are to allow the torn retinaculum to heal and to prevent chronic subluxation. Your doctor may advise you to begin physiotherapy once the cast is removed.

Surgery

Many patients with peroneal tendon subluxation will eventually require surgery, especially when symptoms have not been controlled with nonsurgical measures. The following are different surgical procedures designed to help the peroneal tendons remain in their proper position.

Retinaculum Repair

Retinaculum repair is gaining popularity. This procedure restores the normal anatomy of the retinaculum that normally reinforces the tendon sheath around the peroneal tendons.

In surgery to repair the retinaculum, the surgeon first makes an **incision** along the back and lower edge of the fibula. This lets the surgeon see the spot where the retinaculum is torn.

Incision



The surgeon uses a burr to create a trough along the fibula bone next to the original attachment of the retinaculum. The edge of the retinaculum is then pulled into the trough and **sutured** in place. The skin is closed with stitches.

Sutured

Retinaculum Reattached

Retinaculum
Groove created



Groove Reconstruction

Groove reconstruction is done to deepen the groove so the peroneals stay in place behind the bottom tip of the fibula. In this procedure, the surgeon first makes an **incision** along the back and lower edge of the fibula bone.


Incision



The surgeon cuts a small flap in the bone near the bottom corner of the fibula. The surgeon then carefully folds the flap back, like a hinge. With the hinge held open, the doctor scoops out a small amount of bone under the flap to **deepen**


groove.

Deepen the Groove



The surgeon closes the flap on its hinge and tamps it in place. A screw may be used to hold the flap down.

Closing the Flap




Next, the tendons are returned to their location behind the tip of the fibula. Repair of the retinaculum may also be done with this procedure (see above). The skin is closed and sutured.

Bony Blocks

The purpose of a bony block is to form a barrier that keeps the tendons from slipping out of place. The block is usually formed with bone taken from the lower end of the fibula bone.

To create a bony block, the surgeon opens the skin along the lower edge of the fibula. The surgeon then measures a small area on the back of the fibula, near the lower tip of the bone. A special tool is used to cut this small section of the bone. The cut only goes partway through the bone.



The surgeon slides the small block of bone backward, out of its original spot. The bone may be rotated slightly to create a solid barrier that will help keep the tendons from sliding around the lower edge of the fibula. A screw is inserted into the small block of bone into the fibula. The screw keeps the bony block in its **new location** until it heals.

The surgeon checks the fit to make sure the tendons can glide behind the new block of bone without slipping out. The skin is then closed and sutured.