


Physiotherapy for Ankle Impingement




Welcome to BodyZone Physiotherapy's overview of the anatomy of the ankle.

The ankle joint is formed where the bones of the lower leg, the tibia and the fibula, connect above the anklebone talus. The tibia is the main bone of the lower leg. The fibula is the small, thin bone along the outer edge of the tibia.


The ankle joint is a **hinge** that allows the foot to move up (*dorsiflexion*) and down (*plantarflexion*). The ankle is a **synovial** joint, meaning it is enclosed in a **joint capsule** that contains a lubricant called *synovial fluid*.

Ankle Joint Movement



©MMG 2003


Dorsiflexion



Plantarflexion


Ankle Joint Capsule

Side view



Posterior view


©MMG 2003




Strong **ligaments** surround and support the ankle joint. The ligament that crosses just above the front of the ankle connects the tibia to the fibula is called the *anterior inferior tibiofibular ligament (AITFL)*. The *anterior talofibular (ATFL)* supports the outer edge of the ankle. The ATFL goes from the tip of the fibula and angles forward to connect to the talus.

The talus rests on the the heelbone (the *calcaneus*). The joint formed between these two bones is called the *subtalar joint*. The calcaneus extends backward below the ankle, forming a shelf on which the talus rests.

Two small bony bumps, called **tuberosities**, project from the back of the talus, one on the inside (*medial*) edge and one on the outer (*lateral*) edge.




In some people the lateral tuberosity is not united to the talus. The separate piece of bone is called an **os trigonum**. Separation of the os trigonum from the talus is usually not a fracture. About 15 percent of people have an os trigonum. An os trigonum sometimes causes problems of impingement in the back of the ankle.



Causes

Pinching of tissues in the front of the ankle is called *anterior impingement*. Athletes who have had several mild or one severe sprain are most likely to have anterior impingement. This is especially true for athletes who repeatedly move the ankle upward (dorsiflexion), such as baseball catchers, basketball and football players, and dancers. Over time, irritation along the front edge of the ankle can lead to impingement.




Irritation in the lower edge of the AITFL and the front of the ATFL can thicken these ligaments. The irritated ligaments become vulnerable to getting pinched between the tibia and talus as the foot is dorsiflexed. These ligaments may

to rub on the joint capsule of the ankle. This can inflame the synovial lining of the capsule, a condition called **synovitis**.

A similar problem can happen after an ankle sprain. As the torn or ruptured ligament heals, the body responds by producing much scar tissue along the front and side of the ankle joint. This creates a small mass of tissue called a *meniscoid*. Dorsiflexing the ankle can trap the tissue between the edge of the ankle joint, causing pain, popping, and a feeling that the ankle will give out and not support your body weight.

Over time, damage from past ankle sprains may also lead to the formation of small projections of bone called **bone spurs**. Bone spurs can form along the bottom ledge of the tibia bone or on the upper surface of the talus. As the ankle is dorsiflexed, the bone spurs may begin to jab into the soft tissues along the front edge of the ankle joint, causing **anterior impingement**.





Posterior impingement occurs in the back of the ankle. It is most common in ballet dancers who must continually point their toes, pointing their foot downward into extreme plantarflexion. Other athletes are rarely affected but may have problems if they routinely plantarflex their feet.

The usual cause of posterior impingement is an os trigonum (described earlier). This normal fragment of bone is located between the distal end of the lateral tuberosity from the talus. When an os trigonum is present, it can cause problems, especially among ballet dancers who constantly rise up on their toes into the dance position called *pointe*. Pointe is a position of extreme plantarflexion. As the foot points downward sharply, the os trigonum can get sandwiched between the bottom edge of the tibia and the top surface of the calcaneus (the heelbone). This can **trap** the tissues above and below the os trigonum, leading to symptoms of posterior impingement.

Posterior impingement can also occur in a ballet dancer who has had a previous ankle sprain. Damage from the previous sprain can lead to the formation of scar tissue or bone spurs that can contribute to the impingement.

sprain may create too much instability in the ankle. As the dancer rises up on the toes, the talus may be free to slide slightly. This allows the shelf of the heelbone to come into contact with the back of the tibia, pinching the soft tissue between. Posterior impingement from ankle instability can also happen in other athletes. But this is uncommon, as forceful plantarflexion is rarely required in other sports.

Posterior Impingement



Symptoms



Anterior impingement may feel like ankle pain that continues long after an ankle sprain. The ankle may feel weak and can't be trusted to hold steady during routine activities. When anterior impingement comes from ligament irritation, tissue thickening are usually felt in front and slightly to the side of the ankle. This is the area of the ATFL. The pain is usually felt as the foot is **forced upward** into dorsiflexion. If the ligaments have irritated the synovium of the ankle joint capsule, throbbing pain and swelling from inflammation (synovitis) may also be felt in this area.

Symptoms of posterior impingement include pain behind the heel or deep in the back of the ankle. There is usually tenderness just behind the bottom tip of the fibula, by the outer ankle bone. Pain is usually worse when the foot is **down** into plantarflexion. A painful clicking sensation may also be felt as the foot is twisted in and out.



Diagnosis

The diagnosis of ankle impingement is usually made by examining the ankle. Our physiotherapist will manipulate the ankle to see which movements or positions cause your pain. If anterior impingement is suspected, we may bend your ankle forward or ask you to squat down. To check for posterior impingement, our therapist may push your foot downward or have you push up on your toes. Tenderness can usually be pinpointed over the tissues that are being pinched.

Your physiotherapist BodyZone Physiotherapy may also refer you to a doctor for X-rays or other diagnostics help in accurately assessing your ankle impingement.

Our Treatment

Non-surgical Rehabilitation

Even if you don't require surgery, you may need to follow a program of rehabilitation exercises.

The physiotherapists at BodyZone Physiotherapy can create a program to help you regain ankle function. It is very important that you improve strength and coordination in the ankle.

Initially our physiotherapist will advise you to rest the ankle for a short time to reduce swelling and pain. A special boot or short-leg cast may be recommended to restrict ankle movement for up to four weeks. Mild pain medication and anti-inflammatory medicine, such as ibuprofen, may also be prescribed. An ice pack can also help alleviate swelling and encourage a faster return of normal ankle movement.

Once you begin your BodyZone Physiotherapy rehabilitation program, your recovery may involve doing a series

including stationary cycling, range of motion, and ankle strengthening.

Post-surgical Rehabilitation

After debridement surgery, patients are usually placed in an ankle splint, and begin their recovery by using crutches. The amount of weight put on the foot is gradually increased over a period of approximately one to two weeks. Although the time varies among individuals, our patients generally advance quickly in rehabilitation and are often able to resume normal activity within four to six weeks.

Rehabilitation after excision of the os trigonum is a slower process. We may advise you to attend therapy sessions for three months, with full recovery sometimes taking up to six months. Patients are often kept in the ankle splint for four to six weeks, and crutches are used during this time as the amount of weight borne on the foot is gradually increased.

After removing the stitches and the ankle brace, our patients are often able to begin formal physiotherapy. When you begin your rehabilitation program at BodyZone Physiotherapy, initial treatments begin with gentle range-of-motion exercises for the ankle and toes. The first few physiotherapy treatments are also designed to help control pain and swelling from surgery. Our therapist may use ice, electrical stimulation treatments, massage and other hands-on procedures to reduce muscle spasm and pain.

As the symptoms from surgery begin to ease, our physiotherapist may show you how to do easy ankle motions on a stationary bicycle. After three or four weeks we may advise you to start doing more active ankle exercises. Exercises are designed to improve the strength in the ankle muscles. Our therapist will also help you regain position sense in the ankle joint to improve its stability.

At BodyZone Physiotherapy, our goal is to help you keep your pain under control, improve your range of motion, and maximize strength and control in your ankle. When your recovery is well under way, regular visits to our office will continue to be a resource, but you will be in charge of doing your exercises as part of an ongoing home program.

Physician Review

Your doctor will probably order X-rays if impingement is suspected. X-rays can show if there are bone spurs on the talus. In cases of posterior impingement, an X-ray can show if an os trigonum is present. You may be asked to stand and rise up on your toes during the X-ray. This helps show if impingement is due to bone pinching the soft tissues.


A bone scan may be recommended in select cases, such as when surgery is being considered. In general, MRI scans are helpful for impingement problems, but they may be ordered to check for other ankle problems that could be causing your pain.

If the doctor believes that pinching in the back of the ankle is from an os trigonum, a numbing medication may be injected into this area. If it feels better, the problem is a posterior impingement from the os trigonum. If the pain doesn't go away, the problem could be in the tendon that runs along the inside edge of the os trigonum.

Your doctor may recommend a steroid injection into the painful area. Steroids are strong anti-inflammatory medications. A steroid injection can help relieve irritation and swelling in the soft tissues that are being pinched, reducing their t

Surgery


If nonsurgical treatments do not work, surgery may be recommended. The type of surgery will vary depending on location and cause of ankle impingement.



Debridement is the most common surgery for anterior ankle impingement. Many surgeons prefer to perform this with an **arthroscope**. An arthroscope is a tiny TV camera that can be inserted into a very small incision. It allows the surgeon to see the area where he or she is working on a TV screen.


To begin, two small incisions are made through the skin on each side of the impingement area. The surgeon inserts the arthroscope to see which area of the tendons or joint capsule are irritated and thickened. The arthroscope lets the surgeon determine if a meniscoid lesion (mentioned earlier) is present. A small shaver is used to clear away (*debride*) irritated tissue from the affected ligaments. The surgeon also **debrides** the tissue forming a meniscoid lesion and any areas of the joint capsule that are inflamed. Small forceps may also be used to clear away irritated or inflamed tissue.

Debridement



Small bone spurs on the tibia or talus are **removed**. If the spurs are large, the surgeon may decide to create a new one over or next to the spur. This allows a special instrument, called an *osteotome*, to be inserted. The surgeon uses this to carefully remove these larger bone spurs.


Bone Spur Removal



Before concluding the procedure, a *fluoroscope* is used to check the debridement and to make sure no bony fragments remain. A fluoroscope is a special X-ray machine that allows the surgeon to see a live X-ray picture on a TV screen during surgery. When the surgeon is satisfied that debridement and removal of bone fragments is complete, the skin is sutured together.

Os Trigonum Excision

The goal of an *os trigonum excision* is to carefully remove (*excise*) the os trigonum to alleviate pinching of the tibiotalar ligament or below it. It is standard to use an *open* surgical method which requires a one- to two-inch incision over the outer back of the ankle. An arthroscope is not routinely used for os trigonum excision because there are many nerves and blood vessels in the back of the ankle that could be injured by an arthroscope.



This surgery begins by placing the patient face down on the operating table. The surgeon makes a small **incision** on the lateral side of the back of the ankle, just behind the outer anklebone. A retractor is used to carefully hold the nearby nerves, and blood vessels out of the way. The surgeon locates the os trigonum. A scalpel is usually sufficient to cut the os trigonum. However, if a bony bridge binds the os trigonum to the talus, the surgeon may need to use a chisel or osteotome.

A fluoroscope is used to check for any remaining bony fragments. When the surgeon is satisfied that all bone fragments have been removed, the skin is stitched together. Patients are placed in a special splint designed to protect the ankle and foot from pointing downward.